

IRF3 Recombinant antibody

Cat: B36232S

Company: HaoKebio

Uniprot ID: Q14653

Applications: IHC:1:50-1:500

Organism: Rabbit

IHC-Polymer:1:200-1:2000

Species reactivity: Human Mouse Rat

IHC-TSA:1:250-1:2500

Molecular Weight Calculation: 47 kDa

WB:1:2000-1:10000

Observed Molecular Weight: 50-60 kDa

Background:

The virul-induced expression of interferon(IFN) genes in infected cells implicate in the interplay of several constitutively expressed and virus-activated transcription factors. A family of IFN regulatory factors(IRFs) have been shown to have a role in the transcription of IFN genes as well as IFN-stimulated genes. IRF3 is a novel key transcriptional regulator of type I IFN-dependent immune responses and involves in the innate immune response against DNA and RNA viruses, by binding to the promoters of IFN. It located in the cytoplasm of uninfected cells in an inactive form, and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, could be phosphorylated by IKK and TBK1 kinases. This induces a conformational change, leading to its dimerization, nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAFI), a complex which activates the transcription of the type I IFN and ISG genes.

Synonyms:

2H6, interferon regulatory factor 3, IRF 3, IRF-3

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Cytoplasm,Nucleus

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.


Storage:

Store at -20 °C for one year.


Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0) , Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:



Sample: Mouse testis, 4% PFA 12-24h



Dilution of 1:10000 incubated at room temperature for 1.5 hours.

Source of Reagents:

发表[中文论文]请标注:IRF3(B36232S)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:IRF3(B36232S) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.